

POLAR
PAM
2644

POLARPAM

STATE OF ALASKA
DEPARTMENT OF NATURAL RESOURCES

BOREAL INSTITUTE
DEC 13 1984
LIBRARY

THE CHEENEETNUK LIMESTONE, A NEW EARLY(?) TO MIDDLE DEVONIAN FORMATION
IN THE MCGRATH A-4 AND A-5 QUADRANGLES, WEST-CENTRAL ALASKA

By
R.B. Blodgett and W.G. Gilbert



PROFESSIONAL REPORT 85

Published by
DIVISION OF GEOLOGICAL AND GEOPHYSICAL SURVEYS


ROSS G. SCHAFF
STATE GEOLOGIST

1983

PROFESSIONAL REPORT 85

THE CHEENEETNUK LIMESTONE, A NEW EARLY(?) TO MIDDLE DEVONIAN FORMATION
IN THE MCGRATH A-4 AND A-5 QUADRANGLES, WEST-CENTRAL ALASKA

By
R.B. Blodgett and W.G. Gilbert



058346

FAIRBANKS, ALASKA
1983

BOREAL INSTITUTE

STATE OF ALASKA

Bill Sheffield, *Governor*

Esther C. Wunnicke, *Commissioner, Dept. of Natural Resources*

Ross G. Schaff, *State Geologist*

Cover photograph: View west toward Cheeneetnuk Thrust. Upper plate of thrust consists of the Cheeneetnuk Limestone (dark), which overlies an unnamed dolomite unit (light). Photograph by W.G. Gilbert, 1978.

CONTENTS

	Page
Introduction	1
Cheeneetnuk Limestone	1
Age	1
Paleoecology	2
Regional geologic setting and correlation	3
Acknowledgments	4
References cited	5
Appendix - Description of fossil localities	6

FIGURES

Figure 1. Location of study area in the McGrath A-4 and A-5 Quadrangles, Alaska	2
2. Stratigraphic section that includes the type section of the Cheeneetnuk Limestone	3

PLATE

Plate 1. Geology of the White Mountain - Cheeneetnuk River area, Alaska	Envelope
---	----------

THE CHEENEETNUK LIMESTONE, A NEW EARLY(?) TO MIDDLE DEVONIAN FORMATION IN THE MCGRATH A-4 AND A-5 QUADRANGLES, WEST-CENTRAL ALASKA

By

R.B. Blodgett¹ and W.G. Gilbert²

INTRODUCTION

This report names and describes the Cheeneetnu Limestone, a new lithostratigraphic unit exposed in the McGrath A-4 and A-5 Quadrangles, west-central Alaska (fig. 1). The Cheeneetnu Limestone immediately overlies the uppermost dolomite unit in a stratigraphic column of lower Paleozoic strata exposed in the White Mountain area (Sainsbury, 1965, fig. 2). Sainsbury (1965) described part of the local Ordovician to Devonian section in a 6,000- to 6,500-ft-thick sequence in which he noted only Middle(?) Ordovician and Devonian(?) faunal horizons; he did not establish any formal lithostratigraphic units.

Subsequent field investigations by DGGS (1977-1979) resulted in the first detailed geologic map of the region (Gilbert, 1981). Within the sequence of Paleozoic rocks mapped by Gilbert on the northwest side of the Farewell fault, map unit 'mDlc', a limestone, is equivalent to the Cheeneetnu Limestone. Various taxonomic groups of a well-preserved, rich, diverse, silicified megafauna from the upper part of this unit have been studied by House and Blodgett (1982) and Rigby and Blodgett (1983).

CHEENEETNUK LIMESTONE

The Cheeneetnu Limestone consists of a sequence of well-bedded, dark-gray, argillaceous, micritic limestone. The formation is prominently exposed along the northwest bank of the southwest-flowing Cheeneetnu River, after which it is named. The formation is exposed in a northeast-trending belt across the map area, as shown on plate 1.

The 457-m-thick (1,500 ft) type section of the Cheeneetnu Limestone is located in sections 16 and 21, T. 23 N., R. 32 W., Seward Meridian, McGrath A-5 Quadrangle (fig. 2; pl. 1). The base of the formation is the conformable contact with the underlying unnamed dolomite unit (unit '1Dd' of Gilbert, 1981) exposed in section 21. The top of the unit is exposed at the conformable contact with the overlying unnamed argillite

and chert unit (unit 'uPzac' of Gilbert, 1981) in section 16.

The contact between the underlying unnamed dolomite unit and the Cheeneetnu Limestone is gradational, and dolomitic horizons occur within the lower part of the Cheeneetnu Limestone. The underlying dolomite unit is typically lighter colored (light-gray, pink, or white) and commonly has a sacchroidal texture. The base of the Cheeneetnu Limestone is marked by the first appearance of bedded, dark-gray, micritic limestone. The lower 93 m (300 ft) of the formation is finely laminated and lacks observable macrofossils. The interval between 93 and 278 m (300 and 990 ft) lacks laminations and contains a sparse, nondiverse fauna of favositid corals, dendroid tabulate corals, stromatoporoids, and crinoid ossicles. The upper part of the formation [78 to 457 m (990 to 1,500 ft) above the base] is characterized by several thick beds of the crowded remains of a richly diverse open-marine fauna. These beds, which are separated by even thicker intervals of unfossiliferous micrites, contain abundant silicified remains of brachiopods, rugose and tabulate corals, gastropods, bivalves, rostroconchs, nautiloids, goniatites, tentaculitids, trilobites, ostracodes, crinoid ossicles, sponges, stromatoporoids, and calcareous algae. The contact with the overlying unnamed argillite and chert unit (unit 'Pzac' of Gilbert, 1981) is both abrupt and conformable.

AGE

Only the upper 156 m (510 ft) of the type section of the Cheeneetnu Limestone (localities 79RB6 and 79RB8-79RB11, fig. 2) and several isolated localities within the upper part of the formation yielded age-diagnostic fossils. Dissolution of limestone in acid indicates that conodonts are extremely scarce. Only one age-diagnostic conodont was recovered, *Polygnathus costatus costatus* (localities 79RB8 and 79RB10; identified by N.M. Savage). Within the Eifelian Stage, Klapper and Ziegler (1979, figs. 3-4) showed a global range for *P. costatus costatus*, from the *costatus costatus* Zone into the overlying *australis* Zone. An early Eifelian age is indicated for both localities within the Cheeneetnu Limestone.

Goniatites have been described from localities

¹Department of Geology, Oregon State University, Corvallis, Oregon 97331-5506.

²DGGS, 230 S. Franklin, Juneau, Alaska 99811.




Figure 1. Location of study area in the McGrath A-4 and A-5 Quadrangles, Alaska.

79RB8 and 79RB9 (House and Blodgett, 1982). House identified *Pinacites* sp. juvenile and *Foordites* cf. *platypleura* (Frech) from locality 79RB8, and *Pinacites jugleri* (Roemer), *P.* cf. *jugleri* and *Subanarcestes?* sp. from locality 79RB9. The closest analogues of these fauna are from Europe and North Africa (House and Blodgett, 1982, p. 1873); representatives of the genus *Pinacites* were not previously recognized in the Americas. Although the *Pinacites jugleri* Zone has been assigned to the upper Eifelian Stage, taxa of that zone may range somewhat earlier (House and Blodgett, 1982, p. 1875). This extended range is in accord with Klapper and others (1978) and Chlupac and others (1979), who demonstrate that the base of the *Pinacites jugleri* Zone is near the base of the *Polygnathus costatus costatus* Zone in the Chotec Limestone of Czechoslovakia.

Brachiopods from the Cheeneetnuk Limestone are essentially limited to the upper part of the formation. Although the brachiopod fauna is consistent with an early Eifelian age, it is not age definitive. Brachiopod faunas of early Eifelian age are poorly known from coeval strata of northwestern Canada and the western Cordillera of the United States, to which the Cheeneetnuk Limestone faunas are most closely allied. Rugose corals from the upper part of the formation indicate a probable Eifelian age. Several taxa [*Cystiphyllodes* sp. cf. *C. macrocystis* McLean (locality 79RB4); *Lekanophyllum* sp. cf. *L. mediale* McLean (locality 79RB12); and *Sociophyllum* sp. cf. *S. glomeratum* (Crickmay) (locality 79RB11)] are comparable to species of the Hume Formation (W.A. Oliver, Jr., written

commun., 1980, 1981) of Eifelian age. Corals from the Cheeneetnuk Limestone are listed in Oliver and others (1975, table 14).

Trilobites from the upper part of the formation that were identified by A.R. Ormiston (written commun., 1981) include *Camsellia* n. sp. (localities 79RB6, 79RB20), *Dechenella* cf. *D. setosa* (79WG184), *Basidechenella?* sp. (locality 79WG184), *Fusinipyge?* sp. (locality 79RB4), and *Otarion* sp. (locality 79RB9). Ormiston (written commun., 1981) reported the genus *Camsellia* "...has been known only from the Hume Formation, District of Mackenzie, Northwest Territories. Its presence in these Alaskan collections would tend to strengthen the Hume affinities of other faunal groups from the McGrath Quadrangle collections and would support an Eifelian age."

We conclude that the uppermost part of the Cheeneetnuk Limestone is of early Eifelian (early Middle Devonian) age. The lack of datable fossils in the lower part of the Cheeneetnuk Limestone and in the underlying unnamed dolomite unit make a positive age determination impossible. On the basis of the relative thickness of the Cheeneetnuk Limestone, we suggest that its base extends into the Lower Devonian Series.

PALEOECOLOGY

The basal 93 m (300 ft) of the Cheeneetnuk Limestone is finely laminated and lacks observable megafaunal remains. The fine lamination of probable algal origin, the micritic lithology, and the absence of fossils

and any evidence of bioturbation suggest deposition in restricted, upper intertidal to supratidal quiet water.

The interval from 93 to 278 m (300 to 990 ft) contains a low-diversity fauna of *in-situ* hemispherical favositid and dendroid tabulate corals and *Amphipora* packed biomicrite. The predominance of the sticklike stromatoporoid *Amphipora* is typical of fore-reef, interreef-platform, and back-reef environments (St. Jean, 1971, p. 1402). However, the general absence of more open-marine faunas such as brachiopods and rugose corals suggests that these beds were deposited in stagnant, somewhat hypersaline (lagoonal), shallow-water conditions. The upper 156 m (510 ft) of the formation consists of prominent, thick beds that contain the silicified remains of a rich, diverse, open-marine fauna. The articulated condition of the brachiopod shells and the nonabraded nature of the shelly fossil remains indicate *in-situ* faunas that represent temporary incursions of open-marine conditions into a restricted environment. The presence of calcareous green algae, such as the receptaculitid *Sphaerospongia tessellata* (localities 79RB8 and 79RB9) and the dasyclad genus *Coelotrochium* (localities 79RB4, 79RB6, 79RB8, 79RB9, 79RB11, 79RB12, and 79WG184), indicates these beds were deposited in photic-zone depths.

The fossil-rich beds are separated by much thicker, nonfossiliferous intervals that probably represent continued deposition in a shallow-water, slightly restricted environment, probably under lagoonal conditions.

The sequence of paleoenvironments in the Cheeneetnuke Limestone suggests an upward-deepening section, starting with basal supratidal-intertidal laminates that grade upward into restricted lagoonal micrites. Continuing upward, the lagoonal micrites interfinger with thin tongues of open-marine limy muds that are ultimately replaced by deep-water siliceous muds and oozes of the unnamed argillite and chert unit. Although reef buildups were not observed in the Cheeneetnuke Limestone, their existence in a more seaward position seems probable because it would explain the overall dominance of quiet-water, somewhat restricted (hypersaline-lagoonal) conditions in the formation.

REGIONAL GEOLOGIC SETTING AND CORRELATION

The Cheeneetnuke Limestone forms the uppermost horizon within an Ordovician to Middle Devonian carbonate-platform sequence exposed on the north side of the Farewell fault in the McGrath A-4 and A-5 Quadrangles (Gilbert, 1981). Ordovician to Devonian strata on the southeast side of the Farewell fault are represented primarily by deep-water rocks; the only recognized shallow-water rocks occur at the top of the sequence (Mamet and Plafker, 1982; Gilbert and Bundtzen, 1983; Bundtzen and Gilbert, 1983). Grantz (1966) suggested that 100 km (62 mi) of dextral displacement occurred along the Farewell fault during Late Cretaceous and Cenozoic time. Tentative correlation of offsets of




Figure 2. Stratigraphic section that includes the type section of the Cheeneetnuke Limestone. See plate for location and appendix for sample data.

the truncated trends of shallow-water platform and deeper water rocks across the Farewell fault suggests a right-lateral displacement of about 68 km (43 mi).

The Ordovician to Devonian shallow-water platform rocks extend to the southwest where they terminate in the Sleetmute and Taylor Mountain Quadrangles as the Holitna Group of Cady and others (1955). To the northeast in the Medfra Quadrangle, a similar sequence of Ordovician to Devonian shallow-water platform-carbonate rocks is also present (Patton and others, 1980; Dutro and Patton, 1981).

Much of pre-Cenozoic Alaska, exclusive of the east-central part of the state, may consist of accreted allochthonous terranes and microplates (Berg and others, 1978; Churkin and Carter, 1979; Churkin and Eberlein, 1977; Coney and others, 1980; Jones and others, 1981, 1982). The allochthonous origin of the Nixon Fork terrane possibly is indicated by its abrupt northern boundary (along the Susulatna Lineament) with rocks of the Innoko terrane (Patton, 1978; Patton and Gilbert, 1982), which are characterized by Upper Paleozoic and Mesozoic chert and volcanic rocks, indicative of "...oceanic or island arc affinities" (Patton, 1978, p. B39). However, Patton and Moll (1982, p. 77) indicate that Innoko terrane rocks were derived from the 'Yukon-Koyukuk sea,' a marginal ocean basin of late Paleozoic to early Mesozoic age, thrust southeastward onto Precambrian and early Paleozoic metasedimentary rocks of the Ruby Geanticline. If the latter case is correct, then the Nixon Fork terrane may have originally been contiguous with the Precambrian and early Paleozoic sedimentary rocks of the Ruby Geanticline (Ruby terrane of Coney and others, 1980; Jones and others, 1981).

Some workers believe that the facies difference between the Nixon Fork and Dillinger terranes represents a general increase in water depth to the southeast during early and mid-Paleozoic time (Dutro and Patton, 1981; Churkin and Trexler, 1981; Gilbert and Bundtzen, 1983). Deep-water facies are recognized within a lower Paleozoic sequence described by Churkin and others (1977) in the Terra Cotta Mountains; by Bundtzen and others (1982, 1984), Gilbert and others (1982), and Gilbert and Solie (1983) in the McGrath Quadrangle; by Gilbert (1981) and Bundtzen and Gilbert (1983) in the McGrath and Lime Hills Quadrangles south of the Farewell fault; and by J.M. Hoare (written commun., 1979) along the west bank of the Swift River in the Lime Hills C-7 Quadrangle. The proximity of the Nixon Fork and Dillinger terranes and the presence of alldapic limestones obviously derived from the shallow-water, platform-carbonate environment strongly supports the association of the deep- and shallow-water facies through geologic time. A remarkably similar facies transition is found in the Mackenzie Mountains in northwest Canada (Cecile, 1978).

If the Nixon Fork-Dillinger terrane is truly allochthonous in respect to cratonic North America, its

post-Devonian displacement is minor. This conclusion is based on the very close paleobiogeographic affinities of the Nixon Fork terrane shelly faunas with coeval faunas from the Canadian cordillera, as indicated by the following faunal elements: early Middle Devonian brachiopods from the Cheeneetnuk Limestone; early Middle Devonian corals from the Cheeneetnuk Limestone (W.A. Oliver, Jr., written commun., 1980); early Middle Devonian trilobites from the Cheeneetnuk Limestone (A.R. Ormiston, written commun., 1981); Early Devonian ostracodes from the south flank of Limestone Mountain, Medfra B-4 Quadrangle (J.M. Berdan, written commun., 1981); and early Late Devonian ostracodes from the unnamed Frasnian limestone on the south side of the Farewell fault in the McGrath A-5 Quadrangle (W.K. Braun, written commun., 1982). Blodgett (1983) gives a more detailed discussion of the paleobiogeographic affinities of the Devonian faunas of the Nixon Fork terrane.

Potter and others (1980) indicated that a Late Ordovician brachiopod fauna from the Nixon Fork terrane (White Mountain area, McGrath A-4 Quadrangle) shows closest affinities with a fauna from the upper member of the Jones Ridge Formation of east-central Alaska and Yukon Territory, which was then part of North America (Payne and Allison, 1981). Available paleobiogeographic evidence indicates that the Nixon Fork terrane may represent an *in-situ* miogeoclinal tract that flanked the southeast side of the Ruby Geanticline or a rifted miogeoclinal 'sliver' derived from the vicinity of the Canadian Cordillera.

ACKNOWLEDGMENTS

We are grateful to the following people for their identifications of the fossil fauna and flora of the Cheeneetnuk Limestone: rugose corals, W.A. Oliver, Jr. (U.S. Geological Survey, Washington D.C.); trilobites, A.R. Ormiston (Amoco Research Center, Tulsa, Oklahoma); goniatites, M.R. House (University of Hull, Hull, England); bivalves and rostroconchs, John Pojeta, Jr. (U.S. Geological Survey, Washington D.C.); sponges, J.K. Rigby (Brigham Young University, Provo, Utah); tabulate corals, R.A. Flory (California State University, Chico); dactyloconarids and tentaculitids, Frithjof Lutke (Geologisch-Palaontologisches Institut, Darmstadt, West Germany); conodonts, N.M. Savage (University of Oregon, Eugene); receptaculitids, M.H. Nitecki (Field Museum of Natural History, Chicago, Illinois); and the dasyclad alga genus *Coelotrochium*, Siegfried Rietschel (Landessammlungen für Naturkunde, Karlsruhe, West Germany).

This study was supported by grants from the National Science Foundation, (Grant EAR 79-19993), the Union Oil Company of California Foundation, the Geological Society of America (Grant 2423-79), and by DGGS. We appreciate the thoughtful reviews of A.J. Boucot (Oregon State University, Corvallis), B.K.

Jones (Bear Creek Mining Company, Anchorage, Alaska), and M.W. Henning and J.G. Clough (DGGS).

REFERENCES CITED

- Berg, H.C., Jones, D.L., and Coney, P.J., 1978, Map showing pre-Cenozoic tectonostratigraphic terranes of southeastern Alaska and adjacent area: U.S. Geological Survey Open-file Report 78-1085, scale 1:1,000,000.
- Blodgett, R.B., 1983, Paleobiogeographic affinities of Devonian fossils from the Nixon Fork terrane, southwestern Alaska, in Stevens, C.H., ed., Pre-Jurassic rocks in western North American suspect terranes: Society of Economic Paleontologists and Mineralogists (Pacific Section), p. 125-130.
- Bundtzen, T.K., and Gilbert, W.G., 1983, Outline of geology and mineral resources of upper Kuskokwim region, Alaska, in Proceedings of the 1982 Symposium: Journal of the Alaska Geological Society, v. 3, p. 101-117.
- Bundtzen, T.K., Kline, J.T., and Clough, J.G., 1982, Preliminary geologic map of the McGrath B-2 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Open-file Report 149, 22 p., 1 sheet.
- Bundtzen, T.K., Smith, T.E., Kline, J.T., and Albanese, M.A., 1984, Preliminary geologic map of the McGrath A-2 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Report of Investigations [in press].
- Cady, W.M., Wallace, R.E., Hoare, J.M., and Webber, E.J., 1955, The central Kuskokwim region, Alaska: U.S. Geological Survey Professional Paper 268, 132 p.
- Cecile, M.P., 1978, Report on Road River stratigraphy and the Misty Creek embayment, Bonnet Plume (106B), and surrounding map-areas, Northwest Territories: Geological Survey of Canada Paper 78-2A, p. 371-377.
- Chlupac, Ivo, Lukes, Pavel, and Zikmundova, Jana, 1979, The Lower/Middle Devonian boundary beds in the Barrandian area, Czechoslovakia: *Geologica et Palaeontologica*, v. 13, p. 125-156.
- Churkin, Michael, Jr., and Carter, Claire, 1979, Collision-deformed Paleozoic continental margin in Alaska - A foundation for microplate accretion [abs]: Geological Society of America Abstracts with Programs, v. 11, no. 3, p. 72.
- Churkin, Michael, Jr., and Eberlein, G.D., 1977, Ancient borderland terranes of the North American Cordillera: Correlation and microplate tectonics: Geological Society of America Bulletin, v. 88, p. 769-786.
- Churkin, Michael, Jr., Reed, B.L., Carter, Claire, and Winkler, G.R., 1977, Lower Paleozoic graptolitic section in the Terra Cotta Mountains, southern Alaska Range, in Blean, K.M., ed., The United States Geological Survey in Alaska: Accomplishments during 1976: U.S. Geological Survey Circular 751-B, p. B37-B38.
- Churkin, Michael, Jr., and Trexler, J.H., Jr., 1981, Continental plates and accreted oceanic terranes in the Arctic, in Nairn, A.E.M., Stehli, F.G., and Churkin, Michael, Jr., eds., The ocean basins and margins: The Arctic Ocean, v. 5, p. 1-20.
- Coney, P.J., Jones, D.L., and Monger, J.W.H., 1980, Cordilleran suspect terranes: *Nature*, v. 288, p. 329-333.
- Dutro, J.T., Jr., and Patton, W.W., Jr., 1981, Lower Paleozoic platform sequence in the Medfra Quadrangle, west-central Alaska, in Albert, N.R.D., and Hudson, Travis, eds., The United States Geological Survey in Alaska: Accomplishments during 1979: U.S. Geological Survey Circular 823-B, p. B42-B44.
- Gilbert, W.G., 1981, Preliminary geologic map and geochemical data, Cheeneetnu River area, Alaska: Alaska Division of Geological and Geophysical Surveys Open-file Report 153, 10 p., scale 1:63,360, 2 sheets.
- Gilbert, W.G., and Bundtzen, T.K., 1983, Paleozoic stratigraphy of the Farewell area, southwest Alaska Range, Alaska, in New developments in the Paleozoic geology of Alaska and the Yukon: Alaska Geological Society Symposium, Anchorage, 1983, p. 10-11.
- Gilbert, W.G., and Solie, D.N., 1983, Preliminary bedrock geology of McGrath A-3 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Report of Investigations 83-7, 1 sheet.
- Gilbert, W.G., Solie, D.N., and Dickey, D.B., 1982, Preliminary bedrock geology of the McGrath B-3 Quadrangle, Alaska: Alaska Division of Geological and Geophysical Surveys Open-file Report 148, 1 sheet.
- Grantz, Arthur, 1966, Strike-slip faults in Alaska: U.S. Geological Survey Open-file Report 66-255, 82 p.
- House, M.R., and Blodgett, R.B., 1982, The Devonian goniatite genera *Pinacites* and *Foordites* from Alaska: Canadian Journal of Earth Sciences, v. 19, p. 1873-1876.
- Jones, D.L., Silbering, N.J., Berg, H.C., and Plafker, George, 1981, Map showing tectonostratigraphic terranes of Alaska, columnar sections, and summary description of terranes: U.S. Geological Survey Open-file Report 81-792, 20 p., 2 sheets.
- Jones, D.L., Silbering, N.J., Gilbert, W.G., and Coney, P.N., 1982, Character, distribution, and tectonic significance of accretionary terranes in the central Alaska Range: Journal of Geophysical Research, v. 87, no. B5, p. 3709-3717.
- Klapper, Gilbert, and Ziegler, Willi, 1979, Devonian conodont biostratigraphy, in House, M.R., Scrutton, C.T., and Bassett, M.G., eds., The Devonian System: Special Papers in Palaeontology no. 23, p. 199-224.
- Klapper, Gilbert, Ziegler, Willi, and Mashkova, T.V., 1978, Conodonts and correlation of Lower-Middle Devonian boundary beds in the Barrandian area of Czechoslovakia: *Geologica et Palaeontologica*, v. 12, p. 103-116.
- Mamet, B.L., and Plafker, George, 1982, A Late Devonian (Frasnian) microbiota from the Farewell-Lyman Hills area, west-central Alaska: U.S. Geological Survey Professional Paper 1216-A, 12 p., 4 sheets.

- Oliver, W.A., Jr., Merriam, C.W., and Churkin, Michael, Jr., 1975, Ordovician, Silurian, and Devonian corals of Alaska, in *Paleozoic corals of Alaska*: U.S. Geological Survey Professional Paper 823-B, p. 13-44, pls. 1-25.
- Patton, W.W., Jr., 1978, Juxtaposed continental and oceanic-island arc terranes in the Medfra Quadrangle, west-central Alaska, in Johnson, K.M., ed., *The United States Geological Survey in Alaska: Accomplishments during 1977*: U.S. Geological Survey Circular 772-B, p. B38-B39.
- Patton, W.W., Jr., and Gilbert, W.G., 1982, Tectonics of west-central Alaska and the adjoining Bering Sea region, in *Western Alaska geology and resource potential*: Alaska Geological Society Symposium, Anchorage, 1982, Programs and Abstracts, p. 35.
- Patton, W.W., Jr., and Moll, E.J., 1982, Structural and stratigraphic sections along a transect between the Alaska Range and Norton Sound, in Coonrad, W.L., ed., *The United States Geological Survey in Alaska: Accomplishments during 1980*: U.S. Geological Survey Circular 844, p. 76-78.
- Patton, W.W., Jr., Moll, E.J., Dutro, J.T., Jr., Silberman, M.L., and Chapman, R.M., 1980, Preliminary geologic map of the Medfra Quadrangle, Alaska: U.S. Geological Survey Open-file Report 80-811A.
- Payne, M.W., and Allison, C.W., 1981, Paleozoic continental margin sedimentation in east-central Alaska: *Geology*, v. 9, no. 6, p. 274-279.
- Potter, A.W., Gilbert, W.G., Ormiston, A.R., and Blodgett, R.B., 1980, Middle and Upper Ordovician brachiopods from Alaska and northern California and their paleogeographic implications [abs.]: *Geological Society of America Abstracts with Programs*, v. 12, no. 3, p. 147.
- Rigby, J.K., and Blodgett, R.B., 1983, Early Middle Devonian sponges from the McGrath Quadrangle of west-central Alaska: *Journal of Paleontology*, v. 57, no. 4, p. 773-786.
- Sainsbury, C.L., 1965, Previously undescribed Middle(?) Ordovician, Devonian(?), and Cretaceous(?) rocks, White Mountain area, near McGrath, Alaska, in *Geological Survey Research 1965*: U.S. Geological Survey Professional Paper 525-C, p. C91-C95.
- St. Jean, Joseph, Jr., 1971, Paleobiologic considerations of reef stromatoporoids, in Yochelson, E.L., ed., *North American Paleontological Convention*, Chicago, September 5-7, 1969, v. 2, p. 1389-1429.
- 79RB6 stone scree slope on the west side of the southeast-flowing stream in sec. 22. (USGS 10321-SD) Lat. 62°04'28" N., long. 155°09'21" W. NE1/4 SW1/4 NE1/4 NW1/4 sec. 21, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle), elevation approximately 2,500 ft. Silicified, 1.5-m-thick (5 ft) fossil horizon, 156 m (503 ft) below top of the Cheeneetnuk Limestone.
- 79RB8 (USGS 10061-SD) Lat. 62°04'33" N., long. 155°09'23" W. NE1/4 NW1/4 NE1/4 NW1/4 sec. 21, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Approximately 3.0-m-thick (10 ft) silicified fossil horizon, 102 m (333 ft) below the top of Cheeneetnuk Limestone.
- 79RB9 (USGS 10062-SD) Lat. 62°04'33" N., long. 155°09'23" W. NE1/4 NW1/4 NE1/4 NW1/4 sec. 21, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Approximately 3.0-m-thick (10 ft) silicified fossil horizon 82 m (269 ft) below the top of Cheeneetnuk Limestone.
- 79RB10 (USGS 10094-SD) Lat. 62°04'39" N., long. 155°09'25" W. NE1/4 SW1/4 SE1/4 SW1/4 sec. 16, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Silicified fossil horizon 34 m (113 ft) below top of the Cheeneetnuk Limestone.
- 79RB11 (USGS 10095-SD) Lat. 62°04'50" N., long. 155°09'25" W. NE1/4 SW1/4 SE1/4 SW1/4 sec. 16, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Silicified fossil horizon 25 m (83 ft) below the top of Cheeneetnuk Limestone.
- 79RB12 (USGS 10063-SD) Lat. 62°05'15" N., long. 155°07'27" W. SE1/4 SW1/4 NE1/4 NW1/4 sec. 15, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Rubble crop of silicified fossils just above a band of prominent white dolomite on the south side of an east-dipping slope.
- 79RB20 (USGS 10097-SD) Lat. 62°12'03" N., long. 154°59'43" W. Center of SW1/4 NE1/4 sec. 5, T. 24 N., R. 31 W., Seward Meridian (McGrath A-4 Quadrangle).
- 79WG184 (USGS 10098-SD) Lat. 62°13'02" N., long. 154°53'40" N. SE1/4 SE1/4 NE1/4 NE1/4 sec. 31, T. 25 N., R. 30 W., Seward Meridian (McGrath A-4 Quadrangle).

APPENDIX

DESCRIPTION OF FOSSIL LOCALITIES³

- 79RB4 (USGS 10060-SD) Lat. 62°04'03" N., long. 155°07'05" W. NE1/4 SW1/4 NW1/4 SE1/4 sec. 22, T. 23 N., R. 32 W., Seward Meridian (McGrath A-5 Quadrangle). Silicified fossils recovered from talus of lime-

³U.S. Geological Survey coral collections submitted to W.A. Oliver, Jr., for identification.

STAFF

Ross G. Schaff, State Geologist
W.W. Barnwell, Deputy State Geologist

S.A. Aasland, Clerk-typist	D.L. Krouskop, Geophysicist
D.D. Adams, Geological assistant	G.M. Laird, Cartographer
M.D. Albanese, Geologist	E.M. Lamey, Administrative assistant
R.D. Allely, Hydrologist	F.L. Larson, Publications specialist
M.A. Armstrong, Publications specialist	S.A. Liss, Geological assistant
M.R. Ashwell, Minerals laboratory technician	W.E. Long, Hydrologist
E.E. Becia, Natural resource manager	L.L. Lueck, Geologist
R.M. Bennett, Natural resource manager	W.M. Lyle, Geologist
D.L. Bertossa, Geological assistant	S.F. Mack, Hydrologist
J.E. Bittner, Historian	C.L. Mahan, Accounting technician
J.D. Blum, Geological assistant	R.A. Mann, Clerk
T.K. Bundtzen, Geologist	G.D. March, Geologist
C.D. Burgess, Clerk-typist	M.A. Maurer, Hydrologist
L.E. Burns, Geologist	G.A. McCoy, Hydrologist
S.J. Carrick, Hydrologist	D.L. McGee, Geologist
M.S. Christy, Geologist	R.D. Merritt, Geologist
K.H. Clautice, Geological assistant	J.F. Meyer, Geophysicist
R.A. Clay, Hydrologist	R.J. Michels, Supply technician
J.G. Clough, Geologist	C.W. Mishler, Historian
E.J. Collazzi, Hydrologist	M.A. Moorman, Geologist
R.A. Combellick, Geologist	R.J. Motyka, Geologist
P.L. Coonrod, Secretary	C.G. Mull, Geologist
N.W. Crosby, Operations research analyst	J.A. Munter, Hydrologist
I.E. Cutts, Clerk-typist	D.J. Mursch, Clerk-typist
C.L. Daniels, Publications specialist	J.N. Newgaard, Accounting technician
N.L. Dann, Clerk-typist	C.J. Nye, Geologist
J.N. Davies, Geologist	K.E. Ohlund, Clerk-typist
L.L. Dearborn, Hydrologist	C.D. Olivera, Data entry clerk
D.R. Debenham, Clerk-typist	K.S. Pearson, Cartographer
J.E. Decker, Geologist	G.H. Pessel, Geologist
G. Dickinson, Natural resource officer	M.K. Polly, Minerals laboratory technician
J.T. Dillon, Geologist	M.E. Pritchard, Cartographer
R.G. Dixon, Archaeologist	B. Ranken, Geologist
G.R. Eakins, Geologist	S.E. Rawlinson, Geologist
L. Eggan, Data processing manager	J.W. Reeder, Geologist
G. Finch, Data entry clerk	D.R. Reger, Archaeologist
D.E. Gibson, Archaeologist	R.D. Reger, Geologist
W.G. Gilbert, Geologist	V.L. Reger, Geological assistant
J.E. Giliam, Clerk-typist	R.R. Reifentstuh, Geological assistant
K.M. Goff, Geological assistant	K.M. Rhoades, Analyst programmer
M.R. Granata, Hydrologist	M.S. Robinson, Geologist
J.J. Hansen, Geophysicist	L.C. Schell, Cartographer
E.E. Harris, Geological assistant	R.D. Shaw, Archaeologist
M.W. Henning, Geologist	T.E. Smith, Geologist
C.M. Hernandez, Clerk-typist	T.N. Smith, Geologist
D.R. Hickmott, Geological assistant	M.J. Stroebele, Clerk-typist
C.E. Holmes, Archaeologist	B.A. Syvertson, Clerk-typist
B.S. Hurtig, Analyst programmer	C.A. Ulery, Geological assistant
M.G. Inghram, Hydrologist	R.G. Updike, Geologist
R.W. Ireland, Hydrologist	N.C. Veach, Chemist
S.A. Jacques, Cartographer	J.L. Weir, Clerk-typist
E.L. Jenks, Analyst programmer	S.M. Weum, Geophysicist
T.G. Johnson, Data control specialist	B.A. White, Operations research analyst
J.J. Jurgens, Analyst programmer	L.P. White, Operations research analyst
J.T. Kline, Geologist	M.J. Wibbenmeyer, Natural resource manager
S.L. Klingler, Archaeologist	B.K. Wilson, Geological assistant
R.W. Kornbrath, Geologist	M.A. Wiltse, Chemist
K.J. Krause, Geologist	M.E. Wright, Clerk-typist

Date Due

058346

58346

Pam:551.734: (*49)
BLO

BLODGETT, R.B. and GILBERT, W.G.
The Cheeneetnuk Limestone, a
new Early(?) to Middle Devonian
formati


DATE LOANED	BORROWER'S NAME	DATE DUE

58346

Pam:551.734: (*49)
BLO


BLODGETT, R.B. and GILBERT, W.G.
The Cheeneetnuk Limestone, a
new Early(?) to Middle Devonian
formati

BOREAL INSTITUTE FOR NORTHERN STUDIES, LIBRARY
THE UNIVERSITY OF ALBERTA
EDMONTON, ALBERTA T6G 2E9
CANADA



MAP SYMBOLS

- Contact
- - - High-angle fault—Dashed where approximately located, queried where questionable, dotted where concealed
- Thrust fault, sawtooth on upper plate; dashed where approximately located, queried where questionable
- Anticline
- Strike and dip of beds
- Inclined
- Vertical
- Overturned
- Bearing and plunge of minor fold axis
- Location of Cheeneetnu Limestone type section




Geology by W.G. Gilbert, 1977-78; revised by Christopher Rye (1977), G.M. Laird (1978-79), and V.M. Pardo (1979). Cartography by G.M. Laird.

GEOLOGY OF THE WHITE MOUNTAIN-CHEENEETNUK RIVER AREA, ALASKA

By
R. B. Blodgett and W. G. Gilbert
1983

CORRELATION OF MAP UNITS



Quaternary	CENOZOIC
Tertiary	
Cretaceous	MESOZOIC
Jurassic	
Triassic	
Carboniferous	PALEOZOIC
Devonian	
Silurian	
Ordovician	
Cambrian	

Nixon Fork terrane
Ks

Dillinger terrane
Ks

FAREWELL FAULT

DESCRIPTION OF MAP UNITS
(after Gilbert, 1981)

- QU** SURFICIAL DEPOSITS, UNDIFFERENTIATED—Includes alluvial and colluvial material.
- TS** SANDSTONE—Medium- to thin-bedded, laminated, friable, moderately sorted, medium- to very fine grained carbonaceous micaceous sandstone. Carbonaceous partings between beds and plant remains common, and friable coal beds 0.5-3.0 m (1.6-16 ft) thick occur locally.
- TSS** CONGLOMERATE AND SANDSTONE—Medium- to very thick bedded, light-tan-weathering, poorly sorted, subrounded to rounded granule to pebble quartz conglomerate and poorly sorted, coarse-grained quartz sandstone. Beds are discontinuous, cross-bedded, and contain carbonaceous fragments and partings. Coarse-grained sandstone is interbedded with beds of brown, fine-grained sandstone and siltstone and dark gray carbonaceous shale 5-20 m (16-66 ft) thick. Limestone clasts become more abundant in lower part of unit.
- Ks** PILLOW BASALT—Dark-green pillow basalt interbedded with minor discontinuous beds of dark-brown, gray, and white argillite and chert. Radiolaria identified suggest Triassic age.
- Oi** LIMESTONE—Very thick bedded, dark-gray limestone. Preliminary identification of corals, brachiopods, and foraminifera suggests Late Devonian age.
- Ols** LIMESTONE, SANDSTONE, AND SILTSTONE—Moderately folded, thick-bedded, dark-gray laminated limestone, micaceous sandstone, and siltstone. Stratigraphic position indicates Devonian age.
- St** MICACEOUS SANDSTONE—Brown-weathering, gray, medium- to fine-grained, micaceous calc-sandstone and minor very thick bedded, dark-gray limestone. Graptolites indicate Silurian age.
- Prac** PRE-HOLOCENE ROCKS NORTHWEST OF FAREWELL FAULT
- Ts** GRANITE—Medium- to coarse-grained, equigranular biotite granite that yields potassium-argon age of 39.8 ± 1.2 m.y. (Gilbert, 1981, p. 3).
- Ks** SILTSTONE AND SANDSTONE—Predominantly very well indurated, laminated, gray siltstone and light-tan, very fine to fine-grained quartz sandstone.
- Prac** ARGILLITE AND CHERT—Very well indurated, dark-gray argillite and dark-gray and white banded argillaceous chert.
- St** CHEENEETNUK LIMESTONE—Well-bedded, dark-gray, argillaceous micritic limestone. Abundant brachiopods, gastropods, trilobites, corals, and conodonts indicate Middle and possibly Early Devonian age.
- Ds** DOLOMITE—Very thick bedded, gray, pink, and white, finely laminated, very coarse to fine-grained dolomite. Stratigraphic position indicates Early Devonian age.
- Dsi** ALGAL LIMESTONE—Discontinuous but generally thick-bedded, buff to white stromatopora limestone with subordinate beds of laminated limestone. Preliminary identification of sparse brachiopod and coral faunas indicates Silurian age.
- Oi** LIMESTONE AND ARGILLITE—Thin- to thick-bedded, cherty laminated limestone, algal limestone, limestone breccia, and black argillite. Preliminary identification of abundant brachiopod and conodont faunas indicates Late Ordovician age.
- Ols** ARGILLACEOUS LIMESTONE AND ARGILLITE—Heterogeneous unit of very thin to thin-bedded, cross-bedded argillaceous limestone, thin-bedded laminated limestone, thin- to thick-bedded stromatopora limestone, and thin- to thick-bedded, orange-weathering siltstone and calcarenite. Age assigned to Ordovician because of stratigraphic relationship to limestone and argillite (Oi) unit.
- Om** BANDED MUDSTONE—Very thin to thick-bedded, dark-gray lime mudstone with flattened lime-mud concretions and very thin bedded mudstone and lime mud exhibiting sedimentary bowl- and dome structures. Age uncertain, but assigned to Ordovician because of stratigraphic position.
- Oel** OOLITIC LIMESTONE AND DOLOMITE—Thick-bedded, oolitic pelletal limestone and very thick bedded, white, oolitic-pelletal dolomite. May overlie thin- to thick-bedded, gray laminated limestone in silters along Farewell fault. Age uncertain.

REFERENCES CITED

Gilbert, W.G., 1981. Preliminary geologic map and geochronologic data, Cheeneetnu River area, Alaska: Alaska Division of Geological and Geophysical Surveys Open-File Report 153, 10 p., scale 1:62,360, 2 sheets.

University of Alberta Library



0 1620 0333 6011



Alaska Department of

**NATURAL
RESOURCES**